



TITLE:

Numerical verification method for positivity of solutions to elliptic equations (Numerical Analysis : New Developments for Elucidating Interdisciplinary Problems II)

AUTHOR(S):

Tanaka, Kazuaki; Sekine, Kouta; Oishi, Shin'ichi

---

CITATION:

Tanaka, Kazuaki ...[et al]. Numerical verification method for positivity of solutions to elliptic equations (Numerical Analysis : New Developments for Elucidating Interdisciplinary Problems II). 数理解析研究所講究録 2017, 2037: 125-140

ISSUE DATE:

2017-07

URL:

<http://hdl.handle.net/2433/236865>

RIGHT:

# Numerical verification method for positivity of solutions to elliptic equations

Kazuaki Tanaka<sup>1,\*</sup>, Kouta Sekine<sup>2</sup>, Shin'ichi Oishi<sup>2</sup>

<sup>1</sup>*Graduate School of Fundamental Science and Engineering, Waseda University, Japan*

<sup>2</sup>*Faculty of Science and Engineering, Waseda University, Japan*

**Abstract.** In this paper, we propose a numerical method for verifying the positivity of solutions to semilinear elliptic equations. We provide a sufficient condition for a solution to an elliptic equation to be positive in the domain of the equation, which can be checked numerically without requiring a complicated computation. We present some numerical examples.

**Key words:** computer-assisted proof, elliptic boundary value problem, existence proof, verified numerical computation, verifying positivity

## 1 Introduction

We are concerned with verified numerical computation methods for solutions to the following elliptic problem:

$$\begin{cases} Lu(x) = f(u(x)), & x \in \Omega, \\ u(x) > 0, & x \in \Omega \end{cases} \quad (1a)$$

$$(1b)$$

with an appropriate boundary condition, e.g., the Dirichlet type

$$u(x) = 0, \quad x \in \partial\Omega, \quad (2)$$

where  $\Omega$  is a bounded domain (i.e., an open connected bounded set) in  $\mathbb{R}^n$  ( $n = 1, 2, 3, \dots$ ),  $f: \mathbb{R} \rightarrow \mathbb{R}$  is a given nonlinear function, and  $L$  is a uniformly elliptic self-adjoint operator.

Equation (1a), including the case with (1b), has been widely studied using analytical methods (see, e.g., [7, 2, 14] and the references therein). Moreover, verified numerical computation methods, which originate from [11, 15] and have been further developed by many researchers, in recent years have turned to be effective to obtain, through computer-assistance, existence and multiplicity results for various concrete examples where purely analytical approaches have failed (see, e.g., [12, 13, 17, 18]). These methods enable us to obtain a concrete ball containing exact solutions to (1a), typically in the sense of the norms  $\|\nabla \cdot\|_{L^2(\Omega)}$  and  $\|\cdot\|_{L^\infty(\Omega)}$ . Therefore, such methods have the additional advantage that quantitative information of solutions to a target equation is provided accurately in a strict mathematical sense. However, irrespective of how small the radius of the ball is, the positivity of some solutions is not ensured without additional considerations. For example, in the homogeneous Dirichlet case (2), it is possible for a solution that is verified by such methods not to be positive near the boundary  $\partial\Omega$ .

In this paper, we propose a numerical method for verifying the positivity of solutions to (1a), in order to verify solutions of (1). Theorem 2.2 provides a sufficient condition for positivity, and only requires a simple numerical computation. This is a generalization of [22, Theorem 2] and [23, Theorem 2.2] so that this can be applied to a wider class of elliptic problems. Indeed, Theorem 2.2 (in this paper) works well for verifying positivity of solutions to the stationary problem of the Allen-Cahn equation (see Section 3), whereas the previous theorems in [22, 23] cannot be applied to this problem.

---

E-mail address: \* imahazimari@fuji.waseda.jp

## 2 Verification of positivity

Throughout this paper, we omit the expression “almost everywhere” for Lebesgue measurable functions, for simplicity. For example, we employ the notation  $u > 0$  in the place of  $u(x) > 0$  a.e.  $x \in \Omega$ . Let  $H^1(\Omega)$  be the first order  $L^2$ -Sobolev space on  $\Omega$ . We define  $H_0^1(\Omega) := \{u \in H^1(\Omega) : u = 0 \text{ on } \partial\Omega \text{ in the trace sense}\}$ , and  $H^{-1}(\Omega)$  denotes the dual space of  $H_0^1(\Omega)$  with the usual sup-norm. Moreover, we assume that  $f(u(\cdot)) \in H^{-1}(\Omega)$  for each  $u \in H^1(\Omega)$ , and denote

$$F : \begin{cases} H^1(\Omega) & \rightarrow H^{-1}(\Omega), \\ u & \mapsto f(u(\cdot)). \end{cases} \quad (3)$$

We introduce the following lemma that is required to prove Theorem 2.2.

**Lemma 2.1.** *Let  $u \in H_0^1(\Omega)$  be a weak solution to (1) with the boundary condition (2), such that*

1.  $F(u) \geq 0$  and  $F(u) \neq 0$ ;
2.  $e(u(\cdot)) < \infty$ , where  $e(x) := f(x)x^{-1}$ ,  $x \in \mathbb{R}$ .

Then,

$$\text{ess sup}\{e(u(x)) : x \in \Omega\} \geq \lambda_1, \quad (4)$$

where  $\lambda_1$  is the first eigenvalue of the problem

$$(-L\phi, v)_{L^2(\Omega)} = \lambda(\phi, v)_{L^2(\Omega)}, \quad \forall v \in H_0^1(\Omega); \quad (5)$$

the derivatives on the left side are understood in the sense of distributions.

*Proof.* Let  $\phi_1 \geq 0$  ( $\phi_1 \neq 0$ ) be the first eigenfunction corresponding to  $\lambda_1$  (see, e.g., [4, Theorems 1.2.5 and 1.3.2] for ensuring the nonnegativeness of the first eigenfunction). Since  $L$  is self-adjoint, it follows that

$$(F(u), \phi_1)_{L^2(\Omega)} = \lambda_1(u, \phi_1)_{L^2(\Omega)}.$$

Therefore,

$$\begin{aligned} & (F(u), \phi_1)_{L^2(\Omega)} \\ &= \int_{\Omega} F(u(x)) u(x)^{-1} \{u(x) \phi_1(x)\} dx \\ &\leq \text{ess sup}\{e(u(x)) : x \in \Omega\} (u, \phi_1)_{L^2(\Omega)} \\ &= \lambda_1^{-1} \text{ess sup}\{e(u(x)) : x \in \Omega\} (F(u), \phi_1)_{L^2(\Omega)}. \end{aligned}$$

The positivity of  $(F(u), \phi_1)_{L^2(\Omega)}$  implies (4). □

Using Lemma 2.1, we are able to prove the following theorem, which provides a sufficient condition for the positivity of solutions to (1a).

**Theorem 2.2.** *Let  $u \in C^2(\Omega) \cap C(\overline{\Omega})$  be a function satisfying (1a) (a boundary value condition is not necessary) such that  $\underline{u} \leq u \leq \overline{u}$  in  $\overline{\Omega}$  for  $\underline{u}, \overline{u} \in C(\overline{\Omega})$ . Function  $u$  is always positive in  $\Omega$ , if the following conditions are satisfied:*

1.  $\underline{u}$  is positive in a nonempty subdomain  $\Omega_+ \subset \Omega$  ( $\Omega_+ \neq \emptyset$ );

2.  $\partial\Omega_- \cap \partial\Omega = \emptyset$ , or  $u = 0$  on  $\partial\Omega_- \cap \partial\Omega$  if  $\partial\Omega_- \cap \partial\Omega \neq \emptyset$ , where  $\Omega_-$  is the interior of  $\Omega \setminus \Omega_+$ ;
3. there exists a domain  $\Omega_-^\circ \supset \Omega_-$  ( $\Omega_-^\circ \neq \Omega_-$ ), s.t.,  $f([0, \max\{\bar{u}(x) : x \in \Omega_-^\circ\}]) \geq 0$ ;
4.  $f([\min\{\underline{u}(x) : x \in \Omega_-\}, 0]) \leq 0$  and  $f([\min\{\underline{u}(x) : x \in \Omega_-\}, 0]) < 0$ ;
5. For a domain  $\hat{\Omega}$  such that  $\Omega_- \subset \hat{\Omega} \subset \Omega$ ,  $e([0, -\min\{\underline{u}(x) : x \in \Omega_-\}]) < \lambda_1(\hat{\Omega})$ , where  $e(x) := f(x)x^{-1}$ , and  $\lambda_1(\hat{\Omega})$  is the first eigenvalue of the problem (5) with the notational replacement  $\Omega = \hat{\Omega}$ .

*Proof.* Assume that  $u$  is not always positive in  $\Omega$ , that is, there exists a point  $x \in \Omega_-$  such that  $u(x) \leq 0$ . The strong maximum principle ensures that  $u$  cannot be zero at an interior minimum, i.e., there exists a point  $x \in \Omega_-$  such that  $u(x) < 0$  (the case that  $u \equiv 0$  in  $\Omega_-$  is generally allowed, but this case is also ruled out due to Assumption 3; note that  $(u \geq) \underline{u} > 0$  in  $\Omega_- \setminus \Omega_-$ ). In other words, there exists a nonempty subdomain  $\Omega'_- \subset \Omega_-$  such that  $u < 0$  in  $\Omega'_-$  and  $u = 0$  on  $\partial\Omega'_-$ . Therefore, the restricted function  $v := -u|_{\Omega'_-}$  can be regarded as a solution to

$$\begin{cases} -Lv(x) = f_-(v(x)) \quad (:= -f(-v(x))) & x \in \Omega'_-, \\ v(x) > 0 & x \in \Omega'_-, \\ v(x) = 0 & x \in \partial\Omega'_-. \end{cases}$$

Since  $f_-(v(\cdot)) \geq 0$ ,  $f_-(v(\cdot)) \not\equiv 0$ , and  $f_-(v(\cdot))v(\cdot)^{-1} < \lambda_1(\hat{\Omega}) < \infty$  in  $\Omega'_-$  due to Assumptions 4 and 5, it follows from Lemma 2.1 that

$$\left( \lambda_1(\hat{\Omega}) > \sup_{x \in \Omega'_-} e(-u(x)) = \right) \sup_{x \in \Omega'_-} \frac{f_-(v(x))}{v(x)} \geq \lambda_1(\Omega'_-),$$

where  $\lambda_1(\Omega'_-)$  is the first eigenvalue of (5) with the notational replacement  $\Omega = \Omega'_-$ . Since the inclusion  $\Omega'_- \subset \hat{\Omega}$  ensures that all functions in  $H_0^1(\Omega'_-)$  can be regarded as functions in  $H_0^1(\hat{\Omega})$  by considering the zero extension outside  $\Omega'_-$ , the inequality  $\lambda_1(\Omega'_-) \geq \lambda_1(\hat{\Omega})$  follows. Thus, we have the contradiction that  $\lambda_1(\hat{\Omega}) > \lambda_1(\hat{\Omega})$ .  $\square$

**Remark 2.3.** Assumption 2 in Theorem 2.2 always holds when  $u$  satisfies the homogeneous Dirichlet boundary condition (2).

**Remark 2.4.** We may employ the condition that  $f([0, \max\{\bar{u}(x) : x \in \Omega_-\} + \varepsilon]) \geq 0$  for a positive number  $\varepsilon$ , instead of Assumption 3. Indeed, since  $\bar{u}$  is a continuous function over  $\Omega$ , there exists a domain  $\Omega_-^\circ \supset \Omega_-$  such that  $\max\{\bar{u}(x) : x \in \Omega_-^\circ\} \leq \max\{\bar{u}(x) : x \in \Omega_-\} + \varepsilon$  for any  $\varepsilon > 0$ .

The following corollary is convenient to prove the positivity of a solution  $u$  that is proven to exist in a ball centered around an approximation  $\hat{u}$ . Indeed, in Section 3, we present numerical examples where the positivity of solutions to (1a), that are in balls centered around their approximations, is verified.

**Corollary 2.5.** Let  $u \in C^2(\Omega) \cap C(\bar{\Omega})$  be a function satisfying (1a) (a boundary value condition is not necessary) such that  $|u(x) - \hat{u}(x)| \leq r$  for all  $x \in \Omega$ , for given  $\hat{u} \in C(\bar{\Omega})$  and  $r > 0$ . Function  $u$  is always positive in  $\Omega$ , if the following conditions are satisfied:

1.  $\hat{u} - r$  is positive in a nonempty subdomain  $\Omega_+ \subset \Omega$  ( $\Omega_+ \neq \emptyset$ );
2.  $\partial\Omega_- \cap \partial\Omega = \emptyset$ , or  $u = 0$  on  $\partial\Omega_- \cap \partial\Omega$  if  $\partial\Omega_- \cap \partial\Omega \neq \emptyset$ , where  $\Omega_-$  is the interior of  $\Omega \setminus \Omega_+$ ;
3.  $f([0, 2r + \varepsilon]) \geq 0$  for a positive number  $\varepsilon$ ;

4.  $f([m-r, 0]) \leq 0$  and  $f([m-r, 0)) < 0$ , where  $m := \min\{\hat{u}(x) : x \in \Omega_-\}$ ;
5. For a domain  $\hat{\Omega}$  such that  $\Omega_- \subset \hat{\Omega} \subset \Omega$ ,  $e([0, -m+r]) < \lambda_1(\hat{\Omega})$ , where  $e(x) := f(x)x^{-1}$ , and  $\lambda_1(\hat{\Omega})$  is the first eigenvalue of the problem (5) with the notational replacement  $\Omega = \hat{\Omega}$ .

**Remark 2.6.** Assumption 3 in Corollary 2.5 is simplified owing to the discussion in Remark 2.4.

### 3 Numerical example

In this section, we present numerical examples in which the positivity of solutions to (1a) is verified. All computations were carried out on a computer with Intel Xeon E7-4830 v2  $\times 4$  processors, 2 TB RAM, CentOS 6.6, and MATLAB 2012b. All rounding errors were strictly estimated using the toolboxes of INTLAB version 9 [20] and KV library version 0.4.36 [5] for verified numerical computations. Therefore, the accuracy of all results was mathematically guaranteed. Hereafter,  $\bar{B}(x, r; \|\cdot\|)$  denotes the closed ball whose center is  $x$ , and whose radius is  $r \geq 0$  in the sense of the norm  $\|\cdot\|$ .

For the first example, we select the case in which  $-L = \Delta$  (the usual Laplace operator),  $f(u(\cdot)) = u(\cdot)^p$  and  $F(u) = u^p$  ( $p = 3, 5$ ), and  $\Omega = (0, 1)^2 \subset \mathbb{R}$ , i.e., we consider the problem of finding solutions to

$$\begin{cases} -\Delta u = u^p & \text{in } \Omega, \end{cases} \quad (6a)$$

$$\begin{cases} u > 0 & \text{in } \Omega, \end{cases} \quad (6b)$$

$$\begin{cases} u = 0 & \text{on } \partial\Omega. \end{cases} \quad (6c)$$

We computed approximate solutions  $\hat{u} \in C(\bar{\Omega})$  to

$$\begin{cases} -\Delta u = |u|^{p-1}u & \text{in } \Omega, \\ u = 0 & \text{on } \partial\Omega, \end{cases} \quad (7)$$

which is displayed in Fig. 1, using a Legendre polynomial basis, i.e., we constructed  $\hat{u}$  as

$$\hat{u} = \sum_{i,j=1}^N u_{i,j} \phi_i \phi_j, \quad u_{i,j} \in \mathbb{R}, \quad (8)$$

where each  $\phi_i$  is defined by

$$\phi_n(x) = \frac{1}{n(n+1)} x(1-x) \frac{dP_n}{dx}(x), \quad n = 1, 2, 3, \dots \quad (9)$$

with the Legendre polynomials  $P_n$  defined by

$$P_n = \frac{(-1)^n}{n!} \left( \frac{d}{dx} \right)^n x^n (1-x)^n, \quad n = 0, 1, 2, \dots \quad (10)$$

We proved the existence of solutions  $u$  to (7) in an  $H_0^1$ -ball  $\bar{B}(\hat{u}, r_1; \|\nabla \cdot\|_{L^2(\Omega)})$  and an  $L^\infty$ -ball  $\bar{B}(\hat{u}, r_2; \|\cdot\|_{L^\infty(\Omega)})$  both centered around the approximations  $\hat{u}$ , using Theorem A.1 [17] combined with the method in [24, 8] (see Section A for details). We then verified the positivity of the verified solutions using Corollary 2.5. For the problem (7), Assumptions 2, 3, and 4 in Corollary 2.5 are always satisfied. Therefore, it is only necessary to confirm Assumptions 1 and 5, where  $e(x) = x^{p-1}$  and  $\hat{\Omega} = \Omega$ . Note that the verified solutions have the regularity to be in

$C^2(\Omega) \cap C(\overline{\Omega})$ , regardless of the regularity of the corresponding approximations  $\hat{u}$ . Indeed, for each  $h \in L^2(\Omega)$ , the problem

$$\begin{cases} -\Delta u = h & \text{in } \Omega, \\ u = 0 & \text{on } \partial\Omega \end{cases}$$

has a unique solution  $u \in H^2(\Omega)$ , such as when  $\Omega$  is a bounded convex domain with a piecewise  $C^2$  boundary (see, e.g., [3, Section 3.3]). Therefore, the so-called bootstrap argument ensures that a weak solution  $u \in H_0^1(\Omega)$  to (7) on such a domain  $\Omega$ , is in  $C^\infty(\Omega) (\subset C^2(\Omega))$ . Table. 1 presents the verification result, which ensures the positivity of the verified solutions to (7) centered around  $\hat{u}$ , owing to the condition that  $e([0, -m_{\Omega_-} + r]) \leq (-m_{\Omega} + r)^{p-1} < \lambda_1(\Omega)$ , where we denote  $m_{\Omega_-} = \min\{\hat{u}(x) : x \in \Omega_-\}$  and  $m_{\Omega} = \min\{\hat{u}(x) : x \in \Omega\}$ .




Figure 1: Approximate solutions to (7) on  $\Omega = (0, 1)^2$  for  $p = 3, 5$ .

Table 1: Verification results for (6) on  $\Omega = (0, 1)^2$ , where  $p = 3$  and 5.

$p$	$N$	$r_1$	$r_2$	$m_{\Omega}$	$(-m_{\Omega} + r)^{p-1} \leq$	$\lambda_1(\Omega)$
3	150	6.636469152e-13	4.363745213e-12	0	1.904227228e-23	$(19.7 \leq) 2\pi^2$
5	150	7.088374332e-13	1.724519836e-10	0	8.844489601e-40	"

In the next example, we consider the stationary problem of the Allen-Cahn equation:

$$\begin{cases} -\varepsilon^2 \Delta u = u - u^3 & \text{in } \Omega, \end{cases} \quad (11a)$$

$$\begin{cases} u > 0 & \text{in } \Omega, \end{cases} \quad (11b)$$

$$\begin{cases} u = 0 & \text{on } \partial\Omega, \end{cases} \quad (11c)$$

where  $\varepsilon > 0$ . We again constructed approximate solutions  $\hat{u}$  to (11a) with (11c), i.e., to

$$\begin{cases} -\Delta u = \varepsilon^{-2}(u - u^3) & \text{in } \Omega, \\ u = 0 & \text{on } \partial\Omega, \end{cases} \quad (12)$$

using a Legendre polynomial basis. These solutions are displayed in Fig. 2. Using Theorem A.1, we again verified the existence of solutions to (12) in the balls  $\overline{B}(\hat{u}, r_1; \|\nabla \cdot\|_{L^2(\Omega)})$  and  $\overline{B}(\hat{u}, r_2; \|\cdot\|_{L^\infty(\Omega)})$ , which are also  $C^2$ -regular. For the problem (12), Assumption 2 in Corollary

2.5 is always satisfied as well. Therefore, it is necessary to confirm Assumptions 1, 3, 4, and 5, where  $e(x) = \varepsilon^{-2}(1 - x^2)$ . Note that Assumption 3 is satisfied if  $2r_2 < 1$ , and Assumption 4 is satisfied if  $-m_{\Omega_-} + r_2 (< -m_{\Omega_-} + r_2) < 1$ . We present the verification results for  $\varepsilon = 0.1$ , 0.05, and 0.025 in Table 2, which ensure the positivity of the verified solutions to (12) centered around  $\hat{u}$ , owing to the condition that  $e([0, -m_{\Omega_-} + r_2]) \leq \varepsilon^{-2} < \lambda_1(\hat{\Omega})$ , where we set  $\hat{\Omega} = (0, 1)^2 \setminus [0.009765625, 0.990234375]^2$  and proved  $\Omega_- \subset \hat{\Omega}$  in all the cases. The upper and lower bounds for the first eigenvalue  $\lambda_1(\hat{\Omega})$  were numerically computed with verification using the method in [9, 8] with a piecewise linear finite element basis.




Figure 2: Approximate solutions to (12) on  $\Omega = (0, 1)^2$ .

Table 2: Verification results for (11) on  $\Omega = (0, 1)^2$ .

$\varepsilon$	$N$	$r_1$	$r_2$	$m_{\Omega} \geq$	$\varepsilon^{-2}$	$\lambda_1(\hat{\Omega}) \in$
0.1	60	5.820766091e-15	4.685104029e-14	-2.5439e-3	1.0e+2	$[0.9585, 1.0032]e+5$
0.05	60	9.584445593e-10	3.937471389e-08	-9.9354e-3	4.0e+2	"
0.025	80	1.385525794e-08	2.611277945e-06	-3.0570e-2	1.6e+3	"

## A Verification theorem for elliptic problems

In this section, we apply the method summarized in [16, 17, 18] to a verified numerical computation for solutions to (1a) with (2). Hereafter, we denote  $V = H_0^1(\Omega)$  and  $V^* = H^{-1}(\Omega)$ . We assume that  $F$  defined in (3) is Fréchet differentiable, and there exists  $q \in L^\infty(\Omega)$  such that

$$F'_u u = qu \quad \text{for } u \in V, \quad (13)$$

where  $F'_u$  is the Fréchet derivative of  $F$  at  $\hat{u} \in V$ , i.e., we may regard  $F'_u$  as an  $L^\infty$  function. Indeed, when we select  $F$  as in the examples of Section 3, this is true. We endow  $V$  with inner product

$$(\cdot, \cdot)_V = (\nabla \cdot, \nabla \cdot)_{L^2(\Omega)} + \tau (\cdot, \cdot)_{L^2(\Omega)}$$

and norm  $\|\cdot\|_V := \sqrt{(\cdot, \cdot)_V}$ , where  $\tau$  is a nonnegative number chosen as

$$\tau > -F'_u(x) \quad (\text{a.e. } x \in \Omega). \quad (14)$$

Note that, since the norm  $\|\cdot\|_V$  monotonically increases with  $\tau$ , the usual norm  $\|\nabla \cdot\|_{L^2(\Omega)}$  is bounded by  $\|\cdot\|_V$  for any  $\tau$ ; therefore,  $\overline{B}(\hat{u}, r_1; \|\nabla \cdot\|_{L^2(\Omega)}) \subset \overline{B}(\hat{u}, r_1; \|\cdot\|_V)$  for any  $r_1 \geq 0$  and

$\tau \geq 0$ . The  $L^2$ -inner product and the  $L^2$ -norm are simply denoted by  $(\cdot, \cdot)$  and  $\|\cdot\|$ , respectively, if no confusion arises. By defining  $\mathcal{F} : V \rightarrow V^*$  as

$$\langle \mathcal{F}(u), v \rangle := (\nabla u, \nabla v) - (F(u), v) \quad \text{for } u, v \in V,$$

we first re-write (1a) with (2) as

$$\mathcal{F}(u) = 0 \text{ in } V^*, \quad (15)$$

and discuss the verified numerical computation for (15). In other words, we first consider the existence of a weak solution to (1a) with (2) (a solution to (15) in  $V$ ), and then we discuss its  $H^2$ -regularity if necessary. The norm bound for the embedding  $V \hookrightarrow L^p(\Omega)$  is denoted by  $C_p$ , i.e.,  $C_p$  is a positive number that satisfies

$$\|u\|_{L^p(\Omega)} \leq C_p \|u\|_V \quad \text{for all } u \in V. \quad (16)$$

Since an explicit upper bound for  $C_p$  is important for the verification theory, formulas that give such an upper bound are provided in B.

### A.1 $H_0^1$ error estimation

We use the following verification theorem for obtaining  $H_0^1$  error estimations for solutions to (15).

**Theorem A.1** ([17]). *Let  $\mathcal{F} : V \rightarrow V^*$  be a Fréchet differentiable operator. Suppose that  $\hat{u} \in V$ , and that there exist  $\delta > 0$ ,  $K > 0$ , and a non-decreasing function  $g$  satisfying*

$$\|\mathcal{F}(\hat{u})\|_{V^*} \leq \delta, \quad (17)$$

$$\|u\|_V \leq K \|\mathcal{F}'_{\hat{u}} u\|_{V^*} \quad \text{for all } u \in V, \quad (18)$$

$$\|\mathcal{F}'_{\hat{u}+u} - \mathcal{F}'_{\hat{u}}\|_{B(V, V^*)} \leq g(\|u\|_V) \quad \text{for all } u \in V, \quad (19)$$

and

$$g(t) \rightarrow 0 \quad \text{as } t \rightarrow 0. \quad (20)$$

Moreover, suppose that some  $\alpha > 0$  exists such that

$$\delta \leq \frac{\alpha}{K} - G(\alpha) \quad \text{and} \quad Kg(\alpha) < 1,$$

where  $G(t) := \int_0^t g(s) ds$ . Then, there exists a solution  $u \in V$  to the equation  $\mathcal{F}(u) = 0$  satisfying

$$\|u - \hat{u}\|_V \leq \alpha. \quad (21)$$

Furthermore, the solution is unique under the side condition (21).

Note that, the Fréchet derivative  $\mathcal{F}'_{\hat{u}}$  of  $\mathcal{F}$  at  $\hat{u} \in V$  is given by

$$\langle \mathcal{F}'_{\hat{u}} u, v \rangle = (\nabla u, \nabla v) - (F'_{\hat{u}} u, v) \quad \text{for } u, v \in V.$$

### Residual bound $\delta$

For  $\hat{u} \in V$  that satisfies  $\Delta \hat{u} \in L^2(\Omega)$ , the residual bound  $\delta$  is computed as

$$C_2 \|\Delta \hat{u} + F(\hat{u})\|_{L^2(\Omega)} (\leq \delta);$$

the  $L^2$ -norm can be computed by a numerical integration method with verification.



### Bound $K$ for the operator norm of $\mathcal{F}'_u{}^{-1}$

We compute a bound  $K$  for the operator norm of  $\mathcal{F}'_u{}^{-1}$  by the following theorem, proving simultaneously that this inverse operator exists and is defined on the whole of  $V^*$ .

**Theorem A.2** ([19]). *Let  $\Phi : V \rightarrow V^*$  be the canonical isometric isomorphism, i.e.,  $\Phi$  is given by*

$$\langle \Phi u, v \rangle := (u, v)_V \quad \text{for } u, v \in V.$$

*If the point spectrum of  $\Phi^{-1}\mathcal{F}'_u$  (denoted by  $\sigma_p(\Phi^{-1}\mathcal{F}'_u)$ ) does not contain zero, then the inverse of  $\mathcal{F}'_u$  exists and*

$$\|\mathcal{F}'_u{}^{-1}\|_{B(V^*, V)} \leq \mu_0^{-1}, \quad (22)$$

where

$$\mu_0 = \min \{ |\mu| : \mu \in \sigma_p(\Phi^{-1}\mathcal{F}'_u) \cup \{1\} \}. \quad (23)$$

The eigenvalue problem  $\Phi^{-1}\mathcal{F}'_u u = \mu u$  in  $V$  is equivalent to

$$(\nabla u, \nabla v) - (F'_u u, v) = \mu (u, v)_V \quad \text{for all } v \in V.$$

Since  $\mu = 1$  is already known to be in  $\sigma(\Phi^{-1}\mathcal{F}'_u)$ , it suffices to look for eigenvalues  $\mu \neq 1$ . By setting  $\lambda = (1 - \mu)^{-1}$ , we further transform this eigenvalue problem into

$$\text{Find } u \in V \text{ and } \lambda \in \mathbb{R} \text{ s.t. } (u, v)_V = \lambda ((\tau + F'_u)u, v) \quad \text{for all } v \in V. \quad (24)$$

Owing to (14), (24) is a regular eigenvalue problem, the spectrum of which consists of a sequence  $\{\lambda_k\}_{k=1}^\infty$  of eigenvalues converging to  $+\infty$ . In order to compute  $K$  on the basis of Theorem A.2, we concretely enclose the eigenvalue  $\lambda$  of (24) that minimizes the corresponding absolute value of  $|\mu| (= |1 - \lambda^{-1}|)$ , by considering the following approximate eigenvalue problem

$$\begin{aligned} &\text{Find } u \in V_N \text{ and } \lambda^N \in \mathbb{R} \\ &\text{s.t. } (u_N, v_N)_V = \lambda^N ((\tau + F'_u)u_N, v_N) \quad \text{for all } v_N \in V_N, \end{aligned} \quad (25)$$

where  $V_N$  is a finite-dimensional subspace of  $V$ .

We estimate the error between the  $k$ th eigenvalue  $\lambda_k$  of (24) and the  $k$ th eigenvalue  $\lambda_k^N$  of (25), by considering the weak formulation of the Poisson equation

$$(u, v)_V = (h, v)_{L^2(\Omega)} \quad \text{for all } v \in V \quad (26)$$

for given  $h \in L^2(\Omega)$ ; it is well known that this equation has a unique solution  $u \in V$  for each  $h \in L^2(\Omega)$ . Moreover, we introduce the orthogonal projection  $P_N^\tau : V \rightarrow V_N$  defined by

$$(P_N^\tau u - u, v_N)_V = 0 \quad \text{for all } u \in V \text{ and } v_N \in V_N.$$

The following theorem enables us to estimate the error between  $\lambda_k$  and  $\lambda_k^N$ .

**Theorem A.3** ([24, 8]). *Suppose that there exists a positive number  $C_N^\tau$  such that*

$$\|u_h - P_N^\tau u_h\|_V \leq C_N^\tau \|h\|_{L^2(\Omega)} \quad (27)$$

for any  $h \in L^2(\Omega)$  and the corresponding solution  $u_h \in V$  to (26). Then,

$$\frac{\lambda_k^N}{\lambda_k^N (C_N^\tau)^2 \|\tau + F'_u\|_{L^\infty(\Omega)} + 1} \leq \lambda_k \leq \lambda_k^N,$$

where  $F'_u$  is regarded as an  $L^\infty$  function owing to (13).

The inequality on the right is well known as a Rayleigh-Ritz bound, which is derived from the min-max principle:

$$\lambda_k = \min_{H_k \subset V} \left( \max_{v \in H_k \setminus \{0\}} \frac{\|v\|_V^2}{\|av\|_{L^2(\Omega)}^2} \right) \leq \lambda_k^N,$$

where we set  $a = \sqrt{\tau + F'_u}$  and the minimum is taken over all  $k$ -dimensional subspaces  $H_k$  of  $V$ . Moreover, proofs of the inequality on the left can be found in [24, 8]. Assuming the  $H^2$ -regularity of solutions to (26) (e.g., when  $\Omega$  is convex [3, Section 3.3]), [24, Theorem 4] ensures the left inequality. A more general statement, that does not require the  $H^2$ -regularity, can be found in [8, Theorem 2.1].

**Remark A.4.** When the  $H^2$ -regularity of solutions to (26) is confirmed a priori, e.g., when  $\Omega$  is convex [3, Section 3.3], (27) can be replaced by

$$\|u - P_N^\tau u\|_V \leq C_N^\tau \|\Delta u + \tau u\|_{L^2(\Omega)} \quad \text{for all } u \in H^2(\Omega) \cap V. \quad (28)$$

We can compute an explicit value of  $C_N^0$  with  $\tau = 0$ , for  $V_N$  spanned by a Legendre basis  $\{\phi_i\}_{i=1}^N$  using [6, Theorem 2.3], and may employ  $C_N^\tau = C_N^0 \sqrt{1 + \tau(C_N^0)^2}$ , since

$$\begin{aligned} \|u - P_N^\tau u\|_\tau^2 &\leq \|u - P_N^0 u\|_\tau^2 = \|\nabla(u - P_N^0 u)\|^2 + \tau \|u - P_N^0 u\|^2 \\ &\leq \|\nabla(u - P_N^0 u)\|^2 + \tau(C_N^0)^2 \|\nabla(u - P_N^0 u)\|^2 = (1 + \tau(C_N^0)^2) \|\nabla(u - P_N^0 u)\|^2 \\ &\leq (1 + \tau(C_N^0)^2) (C_N^0)^2 \|\Delta u\|^2 \leq (1 + \tau(C_N^0)^2) (C_N^0)^2 \|\Delta u + \tau u\|^2, \end{aligned}$$

where the last inequality follows from [24, Lemma 1].

### Local lipschitz bound $\mathcal{F}'_u$

A concrete construction of a function  $g$  satisfying (19) and (20) is important for our verification process. For the nonlinearity  $F(u) = au + bu^p$  with  $p \geq 2$  and  $a, b \in L^\infty(\Omega)$  (this form is applicable to the concrete nonlinearities in Section 3), one can employ

$$g(t) = \|b\|_{L^\infty(\Omega)} p(p-1) C_{p+1}^3 K t \left( \|\hat{u}\|_{L^{p+1}(\Omega)} + C_{p+1} t \right)^{p-2},$$

where  $\delta$  and  $K$  are the respective constants in (17) and (18) for  $\hat{u} \in V$ . This selection can be found in [10, Theorem 3.1].

## A.2 $L^\infty$ error estimation

In this subsection, we discuss a method that gives an  $L^\infty$  error bound for a solution to (1a) with (2) from a known  $H_0^1$  error bound, that is, we compute an explicit bound for  $\|u - \hat{u}\|_{L^\infty(\Omega)}$  for a solution  $u \in V$  to (1a) with (2) satisfying

$$\|u - \hat{u}\|_V \leq \rho \quad (29)$$

with  $\rho > 0$  and  $\hat{u} \in V$ . We assume that  $\Omega$  is convex and polygonal to obtain such an error estimation; this condition gives the  $H^2$ -regularity of solutions to (1a) with (2) (and therefore, ensures their boundedness) a priori. More precisely, when  $\Omega$  is a convex polygonal domain, a weak solution  $u \in V$  to (26) and  $h \in L^2(\Omega)$  is  $H^2$ -regular (see, e.g., [3, Section 3.3]). A solution

$u$  satisfying (29) can be written in the form  $u = \hat{u} + \rho\omega$  with some  $\omega \in V$ ,  $\|\omega\|_V \leq 1$ . Moreover,  $\omega$  satisfies

$$\begin{cases} -\Delta\rho\omega = F(\hat{u} + \rho\omega) + \Delta\hat{u} & \text{in } \Omega, \\ \omega = 0 & \text{on } \partial\Omega, \end{cases}$$

and therefore is also  $H^2$ -regular if  $\Delta\hat{u} \in L^2(\Omega)$ . We then use the following theorem to obtain an  $L^\infty$  error estimation.

**Theorem A.5** ([16]). *For all  $u \in H^2(\Omega)$ ,*

$$\|u\|_{L^\infty(\Omega)} \leq c_0\|u\|_{L^2(\Omega)} + c_1\|\nabla u\|_{L^2(\Omega)} + c_2\|u_{xx}\|_{L^2(\Omega)}$$

with

$$c_j = \frac{\gamma_j}{|\bar{\Omega}|} \left[ \max_{x_0 \in \bar{\Omega}} \int_{\bar{\Omega}} |x - x_0|^{2j} dx \right]^{1/2}, \quad (j = 0, 1, 2),$$

where  $u_{xx}$  denotes the Hesse matrix of  $u$ ,  $|\bar{\Omega}|$  is the measure of  $\bar{\Omega}$ , and

$$\gamma_0 = 1, \quad \gamma_1 = 1.1548, \quad \gamma_2 = 0.22361.$$

For  $n = 3$ , other values of  $\gamma_0$ ,  $\gamma_1$ , and  $\gamma_2$  have to be chosen (see [16]).

**Remark A.6.** *The norm of the Hesse matrix of  $u$  is precisely defined by*

$$\|u_{xx}\|_{L^2(\Omega)} = \sqrt{\sum_{i,j=1}^2 \left\| \frac{\partial^2 u}{\partial x_i \partial x_j} \right\|_{L^2(\Omega)}^2}.$$

Moreover, since  $\Omega$  is polygonal,  $\|u_{xx}\|_{L^2(\Omega)} = \|\Delta u\|_{L^2(\Omega)}$  for all  $u \in H^2(\Omega) \cap V$  (see, e.g., [3]).

**Remark A.7.** *Explicit values of each  $c_j$  are provided for some special domains  $\Omega$  in [16, 17]. According to these papers, one can choose, for  $\Omega = (0, 1)^2$ ,*

$$c_0 = \gamma_0, \quad c_1 = \sqrt{\frac{2}{3}}\gamma_1, \quad \text{and} \quad c_2 = \frac{\gamma_3}{3}\sqrt{\frac{28}{5}}.$$

Applying Theorem A.5, we obtain the following corollaries.

**Corollary A.8.** *Let  $u$  be a solution to (7) with  $p \geq 2$ , satisfying (29) for  $\hat{u} \in V$  such that  $\Delta\hat{u} \in L^2(\Omega)$ . Moreover, let  $c_0$ ,  $c_1$ , and  $c_2$  be as in Theorem A.5. Then,*

$$\begin{aligned} \|u - \hat{u}\|_{L^\infty(\Omega)} &\leq c_0 C_2 \rho + c_1 \rho + \\ &c_2 \left( 2^{p-\frac{3}{2}} p \rho C_3 \sqrt{\|\hat{u}\|_{L^{6(p-1)}(\Omega)}^{2(p-1)} + \frac{\rho^{2(p-1)}}{2p-1} C_{6(p-1)}^{2(p-1)}} + \|\Delta\hat{u} + |\hat{u}|^{p-1} \hat{u}\|_{L^2(\Omega)} \right). \end{aligned} \quad (30)$$

*Proof.* Let us denote  $F(u) = |u|^{p-1}u$  in this proof. Due to Theorem A.5, we have

$$\begin{aligned} \|u - \hat{u}\|_{L^\infty(\Omega)} &= \rho \|\omega\|_{L^\infty(\Omega)} \\ &\leq \rho \left( c_0 \|\omega\|_{L^2(\Omega)} + c_1 \|\omega\|_V + c_2 \|\Delta\omega\|_{L^2(\Omega)} \right) \\ &\leq \rho \left( c_0 C_2 + c_1 + c_2 \|\Delta\omega\|_{L^2(\Omega)} \right). \end{aligned}$$

The last term  $\|\Delta\omega\|_{L^2(\Omega)}$  is estimated by

$$\begin{aligned}\rho \|\Delta\omega\|_{L^2(\Omega)} &= \|F(\hat{u} + \rho\omega) + \Delta\hat{u}\|_{L^2(\Omega)} \\ &= \|F(\hat{u} + \rho\omega) - F(\hat{u}) + F(\hat{u}) + \Delta\hat{u}\|_{L^2(\Omega)} \\ &\leq \|F(\hat{u} + \rho\omega) - F(\hat{u})\|_{L^2(\Omega)} + \|\Delta\hat{u} + F(\hat{u})\|_{L^2(\Omega)}.\end{aligned}$$

Since the mean value theorem ensures that

$$\begin{aligned}&\int_{\Omega} (F(\hat{u} + \rho\omega) - F(\hat{u}))^2 dx \\ &= \int_{\Omega} \left( \rho\omega(x) \int_0^1 F'_{\hat{u}+t\rho\omega}(x) dt \right)^2 dx \\ &= \int_{\Omega} \left( \rho\omega(x) \int_0^1 p |\hat{u}(x) + \rho t\omega(x)|^{p-1} dt \right)^2 dx \\ &= p^2 \rho^2 \int_{\Omega} \omega(x)^2 \left( \int_0^1 |\hat{u}(x) + \rho t\omega(x)|^{p-1} dt \right)^2 dx \\ &\leq p^2 \rho^2 \int_{\Omega} \omega(x)^2 \int_0^1 |\hat{u}(x) + \rho t\omega(x)|^{2(p-1)} dt dx \\ &\leq p^2 \rho^2 \|\omega\|_{L^3(\Omega)}^2 \int_0^1 \left\| |\hat{u} + \rho\omega t|^{2(p-1)} \right\|_{L^3(\Omega)} dt \\ &= p^2 \rho^2 \|\omega\|_{L^3(\Omega)}^2 \int_0^1 \|\hat{u} + \rho\omega t\|_{L^{6(p-1)}(\Omega)}^{2(p-1)} dt \\ &\leq p^2 \rho^2 \|\omega\|_{L^3(\Omega)}^2 \int_0^1 \left( \|\hat{u}\|_{L^{6(p-1)}(\Omega)} + t\rho \|\omega\|_{L^{6(p-1)}(\Omega)} \right)^{2(p-1)} dt \\ &\leq 2^{2(p-1)-1} p^2 \rho^2 \|\omega\|_{L^3(\Omega)}^2 \left\{ \|\hat{u}\|_{L^{6(p-1)}(\Omega)}^{2(p-1)} + \int_0^1 \left( t\rho \|\omega\|_{L^{6(p-1)}(\Omega)} \right)^{2(p-1)} dt \right\} \\ &= 2^{2p-3} p^2 \rho^2 \|\omega\|_{L^3(\Omega)}^2 \left( \|\hat{u}\|_{L^{6(p-1)}(\Omega)}^{2(p-1)} + \frac{\rho^{2(p-1)}}{2p-1} \|\omega\|_{L^{6(p-1)}(\Omega)}^{2(p-1)} \right) \\ &\leq 2^{2p-3} p^2 \rho^2 C_3^2 \left( \|\hat{u}\|_{L^{6(p-1)}(\Omega)}^{2(p-1)} + \frac{\rho^{2(p-1)}}{2p-1} C_{6(p-1)}^{2(p-1)} \right).\end{aligned}$$

it follows that

$$\rho \|\Delta\omega\|_{L^2(\Omega)} \leq 2^{p-\frac{3}{2}} p \rho C_3 \sqrt{\|\hat{u}\|_{L^{6(p-1)}(\Omega)}^{2(p-1)} + \frac{\rho^{2(p-1)}}{2p-1} C_{6(p-1)}^{2(p-1)}} + \|\Delta\hat{u} + F(\hat{u})\|_{L^2(\Omega)}.$$

Consequently, the  $L^\infty$  error of  $u$  is estimated as asserted in (30).  $\square$

**Corollary A.9.** *Let  $u$  be a solution to (12) satisfying (29) for  $\hat{u} \in V$  such that  $\Delta\hat{u} \in L^2(\Omega)$ . Moreover, let  $c_0$ ,  $c_1$ , and  $c_2$  be as in Theorem A.5. Then,*

$$\begin{aligned}\|u - \hat{u}\|_{L^\infty(\Omega)} &\leq c_0 C_2 \rho + c_1 \rho + c_2 \left( \rho \varepsilon^{-2} C_3 \left( 1 + 3 \|\hat{u}\|_{L^{12}(\Omega)}^2 \right) \right. \\ &\quad \left. + 3 \rho C_{12} \|\hat{u}\|_{L^{12}(\Omega)} + \rho^2 C_{12}^2 \right) + \|\Delta\hat{u} + \varepsilon^{-2}(\hat{u} - \hat{u}^3)\|_{L^2(\Omega)}.\end{aligned}\tag{31}$$

*Proof.* Let us denote  $F(u) = \varepsilon^{-2}(u - u^3)$  in this proof. Due to Theorem A.5, we have

$$\|u - \hat{u}\|_{L^\infty(\Omega)} = \rho \|\omega\|_{L^\infty(\Omega)}$$

$$\begin{aligned}
&\leq \rho \left( c_0 \|\omega\|_{L^2(\Omega)} + c_1 \|\omega\|_V + c_2 \|\Delta\omega\|_{L^2(\Omega)} \right) \\
&\leq \rho \left( c_0 C_2 + c_1 + c_2 \|\Delta\omega\|_{L^2(\Omega)} \right).
\end{aligned}$$

The last term  $\|\Delta\omega\|_{L^2(\Omega)}$  is estimated by

$$\begin{aligned}
\rho \|\Delta\omega\|_{L^2(\Omega)} &= \|F(\hat{u} + \rho\omega) + \Delta\hat{u}\|_{L^2(\Omega)} \\
&= \|F(\hat{u} + \rho\omega) - F(\hat{u}) + F(\hat{u}) + \Delta\hat{u}\|_{L^2(\Omega)} \\
&\leq \|F(\hat{u} + \rho\omega) - F(\hat{u})\|_{L^2(\Omega)} + \|\Delta\hat{u} + F(\hat{u})\|_{L^2(\Omega)}.
\end{aligned}$$

Since the mean value theorem ensures that

$$\begin{aligned}
&\int_{\Omega} (F(\hat{u} + \rho\omega) - F(\hat{u}))^2 dx \\
&= \int_{\Omega} \left( \rho\omega(x) \int_0^1 F'(\hat{u}(x) + t\rho\omega(x)) dt \right)^2 dx \\
&= \int_{\Omega} \left( \rho\omega(x) \int_0^1 \varepsilon^{-2} \left\{ 1 - 3(\hat{u}(x) + t\rho\omega(x))^2 \right\} dt \right)^2 dx \\
&= \rho^2 \varepsilon^{-4} \int_{\Omega} \omega(x)^2 \left( \int_0^1 \left\{ 1 - 3(\hat{u}(x) + t\rho\omega(x))^2 \right\} dt \right)^2 dx \\
&= \rho^2 \varepsilon^{-4} \int_{\Omega} \omega(x)^2 \left( \int_0^1 (1 - 3\hat{u}(x)^2 - 6t\rho\omega(x)\hat{u}(x) - 3t^2\rho^2\omega(x)^2) dt \right)^2 dx \\
&\leq \rho^2 \varepsilon^{-4} \|\omega\|_{L^3(\Omega)}^2 \left\| \left( \int_0^1 (1 - 3\hat{u}^2 - 6t\rho\omega\hat{u} - 3t^2\rho^2\omega^2) dt \right)^2 \right\|_{L^3(\Omega)} \\
&= \rho^2 \varepsilon^{-4} \|\omega\|_{L^3(\Omega)}^2 \left\| \int_0^1 (1 - 3\hat{u}^2 - 6t\rho\omega\hat{u} - 3t^2\rho^2\omega^2) dt \right\|_{L^6(\Omega)}^2 \\
&\leq \rho^2 \varepsilon^{-4} \|\omega\|_{L^3(\Omega)}^2 \left( 1 + 3 \|\hat{u}^2\|_{L^6(\Omega)} + 3\rho \|\omega\hat{u}\|_{L^6(\Omega)} + \rho^2 \|\omega^2\|_{L^6(\Omega)} \right)^2 \\
&\leq \rho^2 \varepsilon^{-4} \|\omega\|_{L^3(\Omega)}^2 \left( 1 + 3 \|\hat{u}\|_{L^{12}(\Omega)}^2 + 3\rho \|\hat{u}\|_{L^{12}(\Omega)} \|\omega\|_{L^{12}(\Omega)} + \rho^2 \|\omega\|_{L^{12}(\Omega)}^2 \right)^2 \\
&\leq \rho^2 \varepsilon^{-4} C_3^2 \left( 1 + 3 \|\hat{u}\|_{L^{12}(\Omega)}^2 + 3\rho C_{12} \|\hat{u}\|_{L^{12}(\Omega)} + \rho^2 C_{12}^2 \right)^2
\end{aligned}$$

it follows that

$$\begin{aligned}
&\rho \|\Delta\omega\|_{L^2(\Omega)} \\
&\leq \rho \varepsilon^{-2} C_3 \left( 1 + 3 \|\hat{u}\|_{L^{12}(\Omega)}^2 + 3\rho C_{12} \|\hat{u}\|_{L^{12}(\Omega)} + \rho^2 C_{12}^2 \right) + \|\Delta\hat{u} + F(\hat{u})\|_{L^2(\Omega)}.
\end{aligned}$$

Consequently, the  $L^\infty$  error of  $u$  is estimated as asserted in (31).  $\square$

## B Simple bounds for the needed embedding constants

The following theorem provides the best constant in the classical Sobolev inequality with critical exponents.

**Theorem B.1** (T. Aubin [1] and G. Talenti [21]). *Let  $u$  be any function in  $W^{1,q}(\mathbb{R}^n)$  ( $n \geq 2$ ), where  $q$  is any real number such that  $1 < q < n$ . Moreover, set  $p = nq/(n-q)$ . Then,  $u \in L^p(\mathbb{R}^n)$  and*

$$\left( \int_{\mathbb{R}^n} |u(x)|^p dx \right)^{\frac{1}{p}} \leq T_p \left( \int_{\mathbb{R}^n} |\nabla u(x)|_2^q dx \right)^{\frac{1}{q}}$$

holds for

$$T_p = \pi^{-\frac{1}{2}} n^{-\frac{1}{q}} \left( \frac{q-1}{n-q} \right)^{1-\frac{1}{q}} \left\{ \frac{\Gamma(1+\frac{n}{2}) \Gamma(n)}{\Gamma(\frac{n}{q}) \Gamma(1+n-\frac{n}{q})} \right\}^{\frac{1}{n}}, \quad (32)$$

where  $|\nabla u|_2 = ((\partial u/\partial x_1)^2 + (\partial u/\partial x_2)^2 + \cdots + (\partial u/\partial x_n)^2)^{1/2}$ , and  $\Gamma$  denotes the gamma function.

The following corollary, obtained from Theorem B.1, provides a simple bound for the embedding constant from  $H_0^1(\Omega)$  to  $L^p(\Omega)$  for a bounded domain  $\Omega$ .

**Corollary B.2.** *Let  $\Omega \subset \mathbb{R}^n$  ( $n \geq 2$ ) be a bounded domain. Let  $p$  be a real number such that  $p \in (n/(n-1), 2n/(n-2)]$  if  $n \geq 3$  and  $p \in (n/(n-1), \infty)$  if  $n = 2$ . Moreover, set  $q = np/(n+p)$ . Then, (16) holds for*

$$C_p(\Omega) = |\Omega|^{\frac{2-q}{2q}} T_p,$$

where  $T_p$  is the constant in (32).

*Proof.* By zero extension outside  $\Omega$ , we may regard  $u \in H_0^1(\Omega)$  as an element  $u \in W^{1,q}(\mathbb{R}^n)$ ; note that  $q < 2$ . Therefore, from Theorem B.1,

$$\|u\|_{L^p(\Omega)} \leq T_p \left( \int_{\Omega} |\nabla u(x)|_2^q dx \right)^{\frac{1}{q}}. \quad (33)$$

Hölder's inequality gives

$$\begin{aligned} \int_{\Omega} |\nabla u(x)|_2^q dx &\leq \left( \int_{\Omega} |\nabla u(x)|_2^{q \cdot \frac{2}{q}} dx \right)^{\frac{q}{2}} \left( \int_{\Omega} 1^{\frac{2}{2-q}} dx \right)^{\frac{2-q}{2}} \\ &= |\Omega|^{\frac{2-q}{2}} \left( \int_{\Omega} |\nabla u(x)|_2^2 dx \right)^{\frac{q}{2}}, \end{aligned}$$

that is,

$$\left( \int_{\mathbb{R}^n} |\nabla u(x)|_2^q dx \right)^{\frac{1}{q}} \leq |\Omega|^{\frac{2-q}{2q}} \|\nabla u\|_{L^2(\Omega)}, \quad (34)$$

where  $|\Omega|$  is the measure of  $\Omega$ . From (33) and (34), it follows that

$$\|u\|_{L^p(\Omega)} \leq |\Omega|^{\frac{2-q}{2q}} T_p \|\nabla u\|_{L^2(\Omega)}.$$

For any  $\tau \geq 0$ , it is true that  $\|\nabla u\|_{L^2(\Omega)} \leq \|u\|_V$ . □

**Remark B.3.** The case that  $p = 2$  is ruled out in Corollary B.2, but it is well known that

$$\|u\|_{L^2(\Omega)} \leq \frac{1}{\sqrt{\lambda_1 + \tau}} \|u\|_V,$$

where  $\lambda_1$  is the first eigenvalue of the following problem:

$$(\nabla u, \nabla v) = \lambda(u, v) \quad \text{for all } v \in V. \quad (35)$$

Note that, when  $\Omega = (0, 1)^2$ ,  $\lambda_1 = 2\pi^2$ .

The use of the following theorem enables us to obtain an upper bound of the embedding constant when the first eigenvalue  $\lambda_1$  of (35) is concretely estimated. We employ the smaller of the two estimations of  $C_p$  derived from Corollary B.2 and Theorem B.4.

**Theorem B.4** ([18]). Let  $\lambda_1$  denote the first eigenvalue of the problem (35).

a) Let  $n = 2$  and  $p \in [2, \infty)$ . With the largest integer  $\nu$  satisfying  $\nu \leq p/2$ , (16) holds for

$$C_p(\Omega) = \left(\frac{1}{2}\right)^{\frac{1}{2} + \frac{2\nu-3}{p}} \left[\frac{p}{2} \left(\frac{p}{2} - 1\right) \cdots \left(\frac{p}{2} - \nu + 2\right)\right]^{\frac{2}{p}} \left(\lambda_1 + \frac{p}{2}\tau\right)^{-\frac{1}{p}},$$

where  $\frac{p}{2} \left(\frac{p}{2} - 1\right) \cdots \left(\frac{p}{2} - \nu + 2\right) = 1$  if  $\nu = 1$ .

b) Let  $n \geq 3$  and  $p \in [2, 2n/(n-2)]$ . With  $s := n(p^{-1} - 2^{-1} + n^{-1}) \in [0, 1]$ , (16) holds for

$$C_p(\Omega) = \left(\frac{n-1}{\sqrt{n}(n-2)}\right)^{1-s} \left(\frac{s}{s\lambda_1 + \tau}\right)^{\frac{s}{2}}.$$

## Acknowledgments

The authors would like to express their sincere thanks to Professor M. Plum (Institut für Analysis, Karlsruhe Institut für Technologie, Germany) for his comments that helped to correctly describe the verification theory of PDEs in the appendix of this paper. This work was supported by CREST, Japan Science and Technology Agency. The first author (K.T.) is supported by the Waseda Research Institute for Science and Engineering, the Grant-in-Aid for Young Scientists (Early Bird Program). The second author (K.S.) is supported by JSPS KAKENHI Grant Number 16K17651.

## References

- [1] Thierry Aubin. Problèmes isopérimétriques et espaces de Sobolev. *Journal of Differential Geometry*, 11(4):573–598, 1976.
- [2] Haïm Brézis and Louis Nirenberg. Positive solutions of nonlinear elliptic equations involving critical sobolev exponents. *Communications on Pure and Applied Mathematics*, 36(4):437–477, 1983.
- [3] Pierre Grisvard. *Elliptic problems in nonsmooth domains*, volume 69. SIAM, 2011.
- [4] Antoine Henrot. *Extremum problems for eigenvalues of elliptic operators*. Springer Science & Business Media, 2006.
- [5] Masahide Kashiwagi. KV library, 2016. <http://verifiedby.me/kv/>.

- [6] Seiji Kimura and Nobito Yamamoto. On explicit bounds in the error for the  $H_0^1$ -projection into piecewise polynomial spaces. *Bulletin of informatics and cybernetics*, 31(2):109–115, 1999.
- [7] Pierre Louis Lions. On the existence of positive solutions of semilinear elliptic equations. *SIAM review*, 24(4):441–467, 1982.
- [8] Xuefeng Liu. A framework of verified eigenvalue bounds for self-adjoint differential operators. *Applied Mathematics and Computation*, 267:341–355, 2015.
- [9] Xuefeng Liu and Shin’ichi Oishi. Verified eigenvalue evaluation for the laplacian over polygonal domains of arbitrary shape. *SIAM Journal on Numerical Analysis*, 51(3):1634–1654, 2013.
- [10] P.J. McKenna, F. Pacella, M. Plum, and D. Roth. A uniqueness result for a semilinear elliptic problem: A computer-assisted proof. *Journal of Differential Equations*, 247(7):2140–2162, 2009.
- [11] Mitsuhiro T Nakao. A numerical approach to the proof of existence of solutions for elliptic problems. *Japan Journal of Applied Mathematics*, 5(2):313–332, 1988.
- [12] Mitsuhiro T Nakao. Numerical verification methods for solutions of ordinary and partial differential equations. *Numerical Functional Analysis and Optimization*, 22(3-4):321–356, 2001.
- [13] Mitsuhiro T Nakao and Yoshitaka Watanabe. Numerical verification methods for solutions of semilinear elliptic boundary value problems. *Nonlinear Theory and Its Applications, IEICE*, 2(1):2–31, 2011.
- [14] Tiancheng Ouyang and Junping Shi. Exact multiplicity of positive solutions for a class of semilinear problem, II. *Journal of Differential Equations*, 158(1):94–151, 1999.
- [15] Michael Plum. Computer-assisted existence proofs for two-point boundary value problems. *Computing*, 46(1):19–34, 1991.
- [16] Michael Plum. Explicit  $H^2$ -estimates and pointwise bounds for solutions of second-order elliptic boundary value problems. *Journal of Mathematical Analysis and Applications*, 165(1):36–61, 1992.
- [17] Michael Plum. Computer-assisted enclosure methods for elliptic differential equations. *Linear Algebra and its Applications*, 324(1):147–187, 2001.
- [18] Michael Plum. Existence and multiplicity proofs for semilinear elliptic boundary value problems by computer assistance. *Jahresbericht der Deutschen Mathematiker Vereinigung*, 110(1):19–54, 2008.
- [19] Michael Plum. Computer-assisted proofs for semilinear elliptic boundary value problems. *Japan journal of industrial and applied mathematics*, 26(2-3):419–442, 2009.
- [20] Siegfried M Rump. INTLAB - INTerval LABoratory. In Tibor Csendes, editor, *Developments in Reliable Computing*, pages 77–104. Kluwer Academic Publishers, Dordrecht, 1999. <http://www.ti3.tuhh.de/rump/>.
- [21] Giorgio Talenti. Best constant in Sobolev inequality. *Annali di Matematica pura ed Applicata*, 110(1):353–372, 1976.



- [22] Kazuaki Tanaka, Kouta Sekine, Makoto Mizuguchi, and Shin'ichi Oishi. Numerical verification of positiveness for solutions to semilinear elliptic problems. *JSIAM Letters*, 7:73–76, 2015.
- [23] Kazuaki Tanaka, Kouta Sekine, Makoto Mizuguchi, and Shin'ichi Oishi. Sharp numerical inclusion of the best constant for embedding  $H_0^1(\Omega) \hookrightarrow L^p(\Omega)$  on bounded convex domain. *Journal of Computational and Applied Mathematics*, 311:306–313, 2017.
- [24] Kazuaki Tanaka, Akitoshi Takayasu, Xuefeng Liu, and Shin'ichi Oishi. Verified norm estimation for the inverse of linear elliptic operators using eigenvalue evaluation. *Japan Journal of Industrial and Applied Mathematics*, 31(3):665–679, 2014.